

Selected Paper of note: see citations 35, 46, 48, 93,81, 114

PI Frank Gilliland MD PhD

SUMMARY REPORT OF PROGRESS

The Children's Environmental Health Center (CEHC) was established in 1998 to investigate effects of the environment on children's respiratory health, with a focus on asthma and airway disease. In the following sections, we provide an overview of the contributions made by the CEHC. The Center's organization and mechanisms have allowed us to ask broad scientific questions that require an integrated multidisciplinary team to answer efficiently, and at the same time, provide multidisciplinary training for the next generation of environmental health scientists. We focus on how the integrative and collaborative multidisciplinary structure and flexibility of the Center have promoted the development of effective teams of researchers who tackle timely and important scientific problems to efficiently advance environmental health research and translation of research findings into action.

Over the years since the CEHC's founding, we have made major contributions in multiple timely and important scientific and translational arenas, including three areas highlighted in the following sections:

- (1) Identifying clinically significant adverse effects of ambient air pollution on children's respiratory health;
- (2) Demonstrating that common genetic variants are major contributors to children's susceptibility to environmental stressors;
- (3) Defining important pathophysiologic mechanisms for the chronic effects of air pollutants on children's respiratory health; and
- (4) Translating research findings to elevate environmental health effects into an essential element for decision-making in urban planning and economic development in southern California.

Identifying Clinically Significant Adverse Effects of Ambient Air Pollution on Children's Respiratory Health

As described in several high-impact publications, we have observed that children exposed to elevated levels of ambient air pollutants including ozone, PM, NO₂ and fresh traffic-related emissions show clinically important adverse effects on asthma pathogenesis, lung function development, respiratory symptoms and infections.¹⁻¹³ The Center facilitated the multidisciplinary collaborations that led to the development of cutting-edge exposure assessment methodology integrating spatial statistics, home-based measurement of air pollutants, environmental multi-level statistical modeling, geographical information systems (GIS), meteorology, and epidemiology. Several important papers documenting the adverse respiratory health effects of regional pollution relied on data developed by the exposure assessment core from measurements at regional monitoring sites and statistical models developed for these analyses.^{1, 9-10, 14} In order to assess the impact of local exposures due to fresh traffic emissions, new residential and school exposure indices were developed, based on traffic volume, wind speed and direction and mixing layer height, using a GIS system. Using these models, we have demonstrated associations of asthma prevalence and incidence with traffic-modeled pollution and proximity to major roadways.^{3, 6} We have now collected pollution data from a dense network of locations in each of our study communities. Using these measurements and spatial statistical techniques in a Bayesian framework, we have developed new methods for examining effects of traffic-related air pollution.¹⁵ With the development of sophisticated statistical techniques and the advancement of exposure assessment technology (neither of which would have occurred without Center support) we were able to identify independent adverse pulmonary effects from both regional and local pollution.⁵ This observation will clearly have an impact on the policy approaches to dealing with this dual problem.

Demonstrating that Common Genetic Variants are Major Contributors to Children's Susceptibility to Environmental Stressors

The task of decreasing children's susceptibility to air pollution offers an important strategy that can be used to target public health prevention efforts. Emerging research indicates that modifiable antioxidant defenses may be an important response to air pollution. Airborne particulate pollutants, such as DEPs, are thought to exacerbate lung and cardiovascular diseases through induction of oxidative stress.¹⁶ The role of genes involved in oxidative stress produced by xenobiotics (phase II enzymes) has been examined by using GST genotypes, including *GSTM1* and *GSTT1*, which have a null genotype resulting in no protein product, and *GSTP1*, which has a well-studied functional variant (Ile105Val). In individuals with *GSTM1*-null or *GSTP1*-Ile105 genotypes, DEPs enhanced nasal responses to allergen.¹⁷ Compared to subjects with a functional *GSTM1* genotype, *GSTM1*-null subjects had significantly larger increases in IgE levels (146 vs 13.5 U/mL; $P < .01$) and histamine levels (13.9 vs 6.1 nmol/L; $P = .03$) after a DEP plus allergen challenge. The wild-type

GSTP1 genotype was associated with increased IgE levels (149 vs 29.6 U/mL; $P < .01$) and histamine levels (14.5 vs 6.1 nmol/L; $P = .01$) after the same challenge. None of the GSTs modified the response to allergen alone. Common polymorphisms in *GSTM1* and *GSTP1* powerfully modify the adjuvant effect of DEPs on allergic inflammation and identify a large population susceptible to adverse health effects of DEP exposure.

Defining Important Pathophysiologic Mechanisms for the Chronic Effects of Air Pollutants on Children's Respiratory Health

A growing body of evidence indicates that acute and chronic inflammation contributes to the pathogenesis of several common respiratory diseases and conditions.¹⁸⁻²¹ Oxidative and nitrosative stress play important roles in regulating immune responses and subsequent tissue responses. Oxidative/nitrosative stress occurs when oxidant and/or nitrosant burden exceeds the buffering capacity of airway antioxidant/nitrosant defenses.²² In the previous grant period, we investigated whether high ambient air pollutant exposures are associated with airway inflammation as assessed by exhaled nitric oxide (FeNO measured at the conventional 50ml/sec flow rate) in more than 2700 children. We found that exposures to PM_{2.5} mass concentration (seven-day average prior to the test) were associated with increased FeNO. We then investigated whether genetic variants were associated with airway inflammation and children's susceptibility to airway inflammation from ambient air pollution. We found substantial evidence that variation in several genes in the NO synthesis pathway including *NOS2A* and *ARG2* are associated with FeNO, susceptibility to air pollution, risk of new onset asthma and lung function growth. Lastly, we determined that children with chronic airway inflammation as indicated by elevated FeNO are at a greater than two-fold increased risk for new onset asthma. The Center's studies of eNO, along with the studies of genetic variants in oxidative stress and inflammatory pathways provide strong support for the role of oxidative/nitrosative stress in mediating the effects of ambient air pollution.

One approach to reducing the effects of air pollutants such as DEPs is through induction of enzymatic antioxidant defenses, especially in individuals with at-risk genetic variants of key antioxidant enzymes. A prototype for this approach is dietary induction of phase II metabolic enzymes to protect against DEPs. In a proof-of-principle study, dietary sulforaphane, a potent inducer of phase II enzymes, was shown to increase enzyme expression and reduce inflammatory responses.²³ In an in vitro cell system, Wan and Diaz-Sanchez²³ investigated whether sulforaphane stimulated phase II enzyme induction and subsequently reduced the effect of diesel extracts on cytokine production. Sulforaphane increased *GSTM1* and *NQO1* expression, as well as GST activity, while reducing cytokine production. In primary bronchial epithelial cells, sulforaphane also blocked the increased production of interleukin 8, granulocyte-macrophage colony-stimulating factor, and interleukin 1 β from primary human bronchial epithelial cells. DEPs have been shown to increase the production of IgE for *in vitro* cell systems and human nasal challenges. The induction of phase II enzymes in B cells by sulforaphane has been shown to reduce the ability of DEPs to increase IgE production.²³ Dietary sulforaphane also increases phase II enzyme expression in nasal cells. These results suggest that sulforaphane or other compounds that induce phase II enzymes have promise as air pollution chemopreventive agents.²⁴ These chemopreventive effects also might be available through dietary modification. Further research is needed to determine whether the adverse effects of air pollution can be reduced through interventions tailored to individual genetic susceptibility.

Translating Research Findings to Elevate Environmental Health Effects into an Essential Element for Decision-Making in Urban Planning and Economic Development in Southern California

Decisions about urban planning and regional economic development have important long-term implications for children's health, and in general, have not considered the impacts of exposure to environmental air pollution. Major decisions about the urban infrastructure in southern California and the changing global economy present an opportunity for the Center to have lasting impacts on children's health by translating Center findings about the adverse effects of air pollution to decision-makers and stakeholders. The Community Outreach and Translation Core (COTC) has succeeded in discussing environmental health as a central consideration in the debate about alternative development scenarios, creating effective programs for translation of the scientific findings, and providing community members and groups with the knowledge and tools needed to allow these stakeholders to more effectively have their concerns heard. This high-impact effort has the potential to improve health and prevent disease in millions of children for generations to come.

At the inception of our Center in 1998, the issue of air pollution from the ports and goods movement was not a high priority for regulatory agencies, scientists or even many community residents in the harbor area. The volume of cargo containers handled at the Ports of Los Angeles and Long Beach has doubled since that time. Today these adjacent ports are the largest port complex in the United States and the number one single source of air pollution in the region, in part due to lack of emission regulations on foreign-flagged ships and the

use of older diesel trucks to transport containers out of the Ports. Even with the economic slowdown, the Ports continue to anticipate a doubling of Asian imports into the Ports by the year 2030. Plans are underway to expand marine terminals, freeways, rail yards and other transportation infrastructure to accommodate the flow of goods (half of which are destined for other parts of the country). The COTC and Center investigators first became aware of the emerging situation in 2001 by listening to residents of port communities at our NIEHS Town Meeting, co-sponsored by the CEHC. Since 2001, the COTC has worked to integrate concerns about air pollution's adverse health impacts and related health costs into transportation and land-use decisions, where health concerns previously have seldom been a priority. This effort has led to the participation of center investigators Andrea Hricko and Ed Avol in dozens of regional planning, regulatory, port, and transportation committee meetings. The COTC has also educated dozens of community groups about relevant research findings. Our Center has been recognized for its expertise on the environmental health impacts of international trade, ports and goods movement, by appointment of the COTC director to US EPA's National Environmental Justice Advisory Council's "Goods Movement Work Group," and appointment of Center investigators Hricko and Avol to formal stakeholder committees of the Ports and Metropolitan Transportation Authority.

Summary of Key Research Findings from the Previous Cycle

The CEHC has yielded a wealth of data on the acute and chronic respiratory effects of air pollutants; gene-environment interactions; host susceptibility to air pollutants; novel spatial modeling approaches and methods; and exposure assessment methodology. The interdisciplinary nature of this Center has made it possible to develop an integrated program that is synergistic and provides a larger return on investment than would a single investigator model. In this section, we briefly summarize key research findings from the previous cycle.

Health Effects of Air Pollution and Tobacco Smoke Exposures

1. Current levels of air pollution have chronic, adverse effects on lung development in children from the age of 10 to 18 years, leading to clinically significant deficits in attained FEV₁ as children reach adulthood.⁸
2. Respiratory health in children is adversely affected by local exposures to outdoor NO₂ or other freeway-related pollutants.²⁵
3. Residential traffic exposure is associated with deficits in lung function growth.²⁶
4. Residential traffic exposure is associated with prevalent asthma, lifetime asthma and wheezy phenotype.^{3, 27}
5. New onset asthma in primary school children is independently associated with local traffic-related pollution near homes and near schools.²⁸⁻²⁹
6. Markers of traffic-related air pollution are associated with the onset of asthma, providing further evidence that air pollution exposure contributes to new cases of asthma.³⁰⁻³¹
7. On-road commuting exposure to air pollution increases the risk of asthma.³²
8. Maternal and grandmaternal smoking during pregnancy may increase the risk of childhood asthma.³³
9. Regular cigarette smoking increases the risk for new onset asthma among adolescents, especially among those exposed to maternal smoking during the *in utero* period.³⁴

Susceptibility

1. Variation in several genes in the nitric oxide (NO) synthesis and inflammatory pathways, including NOS2A, ARG2, and 5-LO, are associated with eNO, susceptibility to air pollution, and asthma pathogenesis.³⁵⁻³⁸
2. Variants in cytokine genes are associated with exhaled NO and incident asthma.³⁹⁻⁴⁰
3. The tumor necrosis factor (TNF)-308 GG genotype may have a protective role in asthma pathogenesis depending on airway oxidative stress levels.⁴¹
4. In children with at least one copy of the TNF-308 A variant, exposure to two or more household smokers is associated with a two-fold risk of a school absence due to respiratory illness (RI) and a four-fold risk of lower RI-related school absence compared with unexposed children with the TNF-308 GG genotype.⁴²
5. Glutathione-S-transferase M1 and PI (GSTM1 and GSTP1) modify the adjuvant effect of diesel exhaust particles on allergic inflammation.¹⁷
6. Certain variants in intercellular adhesion molecule-1 (ICAM-1) are associated with reduced risk for asthma. Differences in associations of asthma risk with ICAM-1 were found between African-Americans and non-Hispanic and Hispanic whites.⁴³⁻⁴⁴
7. High microsomal epoxide hydrolase (mEH) activity is associated with increased asthma risk and children with both high mEH activity and GSTP1 105Val are at the greatest risk.⁴⁵⁻⁴⁶
8. The clara cell secretory protein gene (CC16) variant allele at position 38 is associated with susceptibility to asthma and wheezing in African-Americans and the risk of early onset asthma among non-Hispanic Whites with a family history of asthma.⁴⁷

9. Genes in glutathione regulation alter the effect of air pollution on lung function growth⁴⁸ and asthma.⁴⁹
10. Dog ownership enhances symptomatic responses to air pollution in children with asthma.⁵⁰
11. Decreased airway flows predict new onset asthma in preadolescent and adolescent children.⁵⁰⁻⁵¹
12. Children with the val105 variant of GSTP1 may be protected from the increased risk of asthma associated with exercise, especially in high ozone communities.⁵²
13. Functional promoter variants in HMOX-1 are associated with a reduced risk for new-onset asthma among non-Hispanic whites; this protective effect is largest in children residing in low-ozone communities.⁵³⁻⁵⁴
14. Genetic variation across the GST mu locus is associated with eight-year lung function growth.⁵⁵
15. Children of mothers who smoked during pregnancy with variation in GSTM2 have lower lung function.⁵⁵
16. Genetic variants in both the promoter and coding regions of the GSTP1 locus may contribute to the occurrence of childhood asthma and wheezing and may increase susceptibility to adverse effects of tobacco-smoke exposure.⁵⁶
17. A positive association is seen between community-level socio-economic position and prevalent and incident childhood asthma, suggesting a significant role for environmental factors in the etiology of this disease.⁵⁷⁻⁵⁸
18. Parental stress increases the risk of childhood wheeze among children with no parental history of asthma, especially among boys.⁵⁹
19. Parental stress and low socioeconomic status increase the risk of childhood wheeze and asthma associated with traffic-related air pollution.⁶⁰

Mechanism

1. Exhaled NO is a useful biomarker for airway inflammation in large population-based studies.⁶¹⁻⁶³
2. Exposure to particulate matter less than 2.5 microns in diameter (PM_{2.5}) mass concentration (two to seven-day average prior to the test) is associated with increased exhaled NO.⁶⁴
3. Children with chronic airway inflammation, as indicated by elevated eNO, are at increased risk for new-onset asthma.⁶⁵
4. Primary allergic sensitization may be prevented by initial high levels of respiratory allergen exposure through induction of a modified, nonallergic immune response.⁶⁶
5. Evidence from controlled chamber experiments shows that secondhand smoke can exacerbate allergic responses in humans.⁶⁷
6. A natural protective mechanism in B cells from oxidant pollutants, such as diesel particles, is the expression of phase II enzymes through induction of antioxidant response elements.²³
7. Induction of phase II enzymes by the chemical sulforaphane can block DEP-induced enhanced IgE production in B cells and DEP-induced proinflammatory cytokine production in epithelial cells.⁶⁸
8. Pretreatment with sulforaphane inhibits diesel-induced production of IL-8, granulocyte-macrophage colony-stimulating factor, and IL-1beta from primary human bronchial epithelial cells, demonstrating that sulforaphane can mitigate the effect of diesel in respiratory epithelial cells.²⁴
9. A placebo-controlled dose escalation trial demonstrates that sulforaphane from broccoli sprouts induces a potent increase in antioxidant Phase II enzymes in airway cells and suggests enhancement of Phase II enzyme expression as a novel therapeutic strategy for oxidant induced airway disease.⁶⁹

New Methodology

1. Offline eNO field measurements can predict online eNO with concurrent ambient NO measurements.⁷⁰⁻⁷²
2. Field-based extended NO testing of children yields useful information about NO at different levels of the respiratory tract, not obtainable from conventional eNO measurement⁷³
3. The development of models to assess air pollution exposures within cities for assignment to subjects in health studies augments the field of exposure assessment and may help to reduce scientific uncertainties that now impede policy intervention aimed at protecting public health.⁷⁴
4. Methods were developed to optimally locate a dense network of monitoring stations representing land use, transportation infrastructure, and the distribution of at-risk populations; this has widespread applicability for the design of pollution monitoring networks, particularly for measuring traffic pollutants with fine-scale spatial variability.⁷⁵
5. Residential ozone concentrations may be over- or underestimated by measurements at a community monitor, depending on the variation in local traffic in the community.⁷⁶
6. Statistical methods have been developed related to the general multi-level modeling paradigm, with a focus on flexible modeling techniques for nonlinear lung function trajectories in children and their relationship to air pollution. The models address many issues, including ecologic inference, multi-pollutant and subgroup analysis, simultaneous modeling of several outcomes, and exposure measurement error.⁷⁷

7. A testing strategy called the "focused interaction testing framework" (FITF) was developed to identify susceptibility genes involved in epistatic interactions for case-control studies of candidate genes. In CHS data, FITF identified a significant multilocus effect between NQO1, MPO, and CAT, three genes involved in the oxidative stress pathway. In an independent data set, these three genes also show a significant association with asthma status.⁷⁸
8. For analysis of a single candidate gene, the SNP interaction model with phase information (SIMPlE) model can be used to identify important SNPs and underlying haplotype structures across a variety of causal models and genetic architectures.⁷⁹
9. A two-step approach was developed to test association of disease with multiple single nucleotide polymorphisms (SNPs) within a candidate locus. The first step uses principal components (PCs) analysis to compute combinations of SNPs that capture the underlying correlation structure within the locus. The second step uses the PCs directly in a test of disease association.⁸⁰
10. A two-step analysis of genome-wide association study data aimed at identifying genes involved in a gene-environment interaction was developed. The procedure complements standard screening for marginal genetic effects and has the potential to uncover novel genetic signals.⁸¹

Community Outreach and Translation

1. Traditional approaches to the calculation of attributable risk may underestimate the health impact of long-term environmental or other exposures that produce both chronic and acute disease.⁸²
2. Community-based quantitative risk analyses can improve our understanding of health problems and help promote public health in transportation planning.⁸³⁻⁸⁴
3. As ports and goods movement activity expands throughout the United States, outreach efforts in the Center have made health and community impacts a more central part of policy discussions.⁸⁵⁻⁸⁸
4. The large population of children exposed to high levels of outdoor air pollutants and the substantial risks for adverse health effects presents unexploited opportunities to reduce the burden of asthma.⁸⁹

BRIEF PROGRESS SUMMARIES PROJECTS

This Children's Environmental Health Center renewal supports a program of ongoing research examining the genetic and environmental determinants of respiratory diseases. Here we briefly highlight some of the most significant results from the previous five years of Projects 1 (McConnell) and 3 (Gilliland) that will continue.

Urban Air Pollution and Persistent Early Life

In the last five years, we have evaluated three hypotheses about traffic-related air pollution: 1) Prevalent asthma with early onset (in the first 5 years of life) is strongly associated with early life traffic within 100 m of the child's home; 2) Early onset asthma is independently associated with variation of NO, NO₂ and ozone within communities, which represent regional products of atmospheric photochemical oxidation of traffic related pollutants, but which also vary locally with traffic; 3) The effects of air pollutants on asthma risk in children are modified by polymorphisms in GSTM1, GSTP1, NQO1, HO-1, and TNF α , genes involved in response to oxidant air pollutants. We found that early life asthma that was symptomatic during the year prior to study entry at age 5-7 was associated with traffic proximity and with NO, NO₂ and NO_x estimated from dispersion models and traffic volume. There was a dose response relationship between distance to traffic corridors, with increased risk beginning approximately 200 m from a major road and increasing approximately linearly with proximity. The odds ratio was 1.6 for asthma within 75 m of major roads (including freeways, highways and major arterials) where 15% of all participants lived. Risk was larger in children with early life (and longer duration) exposure based on residence at the same address since age 2. Susceptible groups were identified, including increased risk in girls, consistent with previous literature. Risk of asthma symptoms, severity and prevalence (or asthma incidence in independently funded related work) associated with traffic-related local or regional air pollution was found to be modified by genetic variants involved in oxidative/nitrosative stress pathways, including GSTM1 and P1, TNF α , and EPHX1. The associations of lifetime asthma with NO_x, NO₂ and ozone measured in a case-control study were not significant. However, based on these and additional measurements at a total of over 1000 locations across southern California improved models predicting spatial surfaces of these pollutants in our study communities based on traffic distance, volume, meteorology, population density, and other nearby land use. Predicted exposure to NO_x was associated with asthma and strongly associated with novel wheezy phenotype based on number and severity of symptoms. Regional pollution was not associated with asthma across study communities, after adjusting for local traffic-related exposure, although ozone was found to modify the effect of genetic variants on the risk of asthma and asthmatic bronchitis. This study was conducted as a community based participatory research (CBPR) project

in close coordination with COTC goals to develop community capacity to understand and apply research results to policy related prevention. In this context novel risk assessment methods were developed that identified a several-fold increase in air pollution-related emergency room visits and other severe asthma morbidity likely to result from increased asthma rates due to traffic proximity in the study communities of Long Beach and Riverside. Work with community partners to develop estimates of traffic volume on major traffic corridors has led to novel pilot work by Center investigator Scott Fruin to use these methods to improve our prediction model surfaces. CBPR is not explicitly a feature of the renewal application, as we have developed a new risk assessment initiative and a large pending grant application with the Air Quality Management District, our governmental partner in the last cycle of the CEHC. The renewal application is focused on developing a comprehensive estimate of the burden of disease surrounding major traffic corridors in LA County, including the 710 corridor that is the “laboratory” for COTC. Our community engagement “essential characteristic” for the proposal draws upon the strong community partnerships developed in our previous grant cycles, which has led to leveraging of funding by the COTC to establish a community/academic collaborative involving our two CBPR partner organizations and several new partners, described in more detail in the COTC section of this proposal.

Air Pollution, Exhaled Breath Markers and Asthma in Susceptible Children (now Project 2: PM, Extended eNO, Susceptibility, and Lung Development)

In the past five years, we assessed three hypotheses based on a biological model for the oxidative stress and inflammatory pathways involved in the pathogenesis of adverse effects of air pollution: (1) High ambient air pollution exposure is associated with chronic airway inflammation in children as indicated by elevated exhaled NO (eNO), a marker of airway inflammation and oxidative/nitrosative stress; (2) Children’s susceptibility to airway inflammation and oxidative/nitrosative stress from ambient air pollution varies by NOS1, NOS2 and NOS3, GSTM1, GSTP1, NQO1, and HO-1 genotypes; and (3) Children with chronic airway inflammation as indicated by elevated eNO are at increased risk for new onset asthma. We first investigated whether high ambient air pollutant exposures were associated with elevated eNO (marker of inflammation) in more than 2700 children. We found that exposures to particulate matter less than 2.5 microns in diameter (PM_{2.5}) mass concentration (two to seven-day average prior to the test) were associated with increased eNO. In order to investigate the second hypothesis, we assessed whether specific genetic variants were associated with airway inflammation (elevated eNO) and children’s susceptibility to airway inflammation from ambient air pollution. We found strong evidence that variation in several genes on the nitric oxide (NO) synthesis pathway, including NOS2A and ARG2, are associated with eNO, susceptibility to air pollution, risk of new onset asthma and lung function growth. Finally, in order to evaluate hypothesis 3, we investigated whether elevated exhaled NO at baseline is associated with new onset asthma during the follow-up period. We determined that children with chronic airway inflammation as indicated by elevated eNO are at increased risk for new onset asthma. Taken together, these findings support the hypothesis that ambient PM produces inflammation and oxidative/nitrosative stress in the airways and provide evidence that airway inflammation/nitrosative stress contributes to asthma pathogenesis and abnormal lung function development. Furthermore, these findings indicate that variants in genes involved in NO production pathways may affect susceptibility to the adverse effects of air pollution. Based on preliminary data, studies of genes and pathways involved in regulating inflammatory responses are likely to provide crucial information needed to understand the role of the interrelated processes of oxidative/nitrosative stress on inflammation in respiratory disease pathogenesis. The renewal application proposes to build on the unique resources of the CEHC by taking the next steps in filling important gaps in the knowledge base.

: Pollution-Enhanced Allergic Inflammation and Phase II Enzymes

Aim #1: We will test the hypothesis that Phase II enzyme expression in the upper airways are induced by oxidant pollutants and differ between children and adults. We had previously shown that challenge of individuals with diesel exhaust particles (DEP) induced GSTM1 expression in adults.

We expanded these studies and completed similar challenges in 20 children. In adults, DEP induced expression of GSTP1 in a dose-dependent fashion. With increasing concentrations of DEP challenge, there




Figure 2 GSTP1 response to DEP in adults vs. children




Figure 3 Cellular response to DEP in adults vs. children

were higher levels of GSTP1-relative gene expression. Cellular infiltration was inversely correlated with GSTP1 enzyme gene expression. Increased levels of GSTP1 were associated with lower cell count numbers. In children, the degree of cellular infiltration also correlated with dose of nasally administered DEP challenge. Higher concentrations of DEP elicited a larger number of total cells recovered from nasal lavage fluid obtained 24 hours after DEP exposure. However, as compared to adults, this effect in children was more robust at the DEP challenge of 300 μ g. In these children, DEP also induced expression of GSTP1. At DEP concentrations of 30 and 100 μ g, there were higher levels of GSTP1-relative gene expression. However, *GSTP1-relative gene expression for the DEP nasal challenge of 300 μ g was decreased in children as compared to adults.* Similar results were seen with the other three measured Phase II enzymes.

Aim#3: We will determine the role of Phase II enzymes in regulating the adjuvant effects of PM.

We have previously shown that enhancement of Phase II enzyme expression could inhibit enhancement of IgE production by peripheral blood cells *in vitro*. In those experiments, we used sulforaphane—a potent inducer of Phase II enzymes found in cruciferous vegetables. We tested whether this same reagent could block DEP-mediated adjuvant events *in vivo* in a murine model. Groups of six female BALB/c mice matched for age and weight (10-12 weeks) received aerosolized exposures to either saline, ovalbumin (OVA) (1% 20 min), DEP (1 hr) or OVA followed immediately by DEP for ten consecutive days. Mice were administered either vehicle (corn oil) or sulforaphane (4.5 μ mol/mouse/day) by gavage (0.2 ml) for one week prior to the commencement of exposures up until two days after the last exposure. As expected, in the untreated group, IgG1 levels were significantly higher in mice who received both DEP and OVA than in those who were exposed to OVA alone. However, administration of sulforaphane completely ablated the adjuvant effects of DEP.

We then examined the responses of the airway epithelial cell line BEAS-2B and primary normal human bronchial epithelial cells upon treatment with sulforaphane and Phase II enzymes followed by stimulation with diesel extract (0-25 μ g/ml). As expected, sulforaphane upregulated the expression of endogenous antioxidant enzymes in bronchial epithelial cells. Whereas diesel extract stimulated the production of IL-8, GM-CSF, and IL-1 β from normal human bronchial epithelial cells, pre-treatment with sulforaphane for 24 hours inhibited diesel-induced cytokine production in a dose-dependent fashion. These studies suggest that sulforaphane treatment can prevent the production of pro-inflammatory cytokines in respiratory epithelial cells *in vitro*.

Previously, we reported that individuals who lack the ability to make the Phase II enzyme GSTM1 are at increased risk for the pro-inflammatory effects of DEP, and we have shown that enhancement of Phase II enzymes with sulforaphane can inhibit the production of pro-inflammatory cytokines in respiratory epithelial cells *in vitro*. In order to determine whether GSTM1 itself is important in the regulation of inflammatory response to pollutants, we used siRNA to “knock down” the GSTM1 gene in bronchial epithelial cells. Expression could be reduced by more than 90% using this methodology. Knockdown of GSTM1 augmented DEP induced cytokine production in these cells. Thus, IL-8 levels were almost three-fold higher in cells where GSTM1 expression was reduced, compared to sham-treated cells.

Finally, in a placebo-controlled dose escalation trial to investigate the *in vivo* effects of sulforaphane, we demonstrated that consumption of oral sulforaphane contained in broccoli sprouts homogenate (BSH) can

induce a potent increase in antioxidant Phase II enzymes in airway cells. RNA expression for GSTM1, GSTP1, HO-1 and NQO1 was measured in nasal lavage cells by RT-PCR before and after sulforaphane dosing. Increased Phase II enzyme expression occurred in a dose-dependent manner with maximal enzyme induction observed at the highest dose of 200 g broccoli sprouts prepared as BSH. Significant increases were seen in all sentinel Phase II enzymes RNA expression compared to baseline. Phase II enzyme induction was not seen with ingestion of non-sulforaphane containing alfalfa sprouts. These findings suggest enhancement of Phase II enzyme expression as a novel therapeutic strategy for oxidant-induced airway disease.⁶⁹

Our results show that Phase II enzymes can be induced by our model air pollutant, DEP, and are critical in regulating responses and determining susceptibility to these xenobiotics. A principal finding of our results is the discovery that children have enhanced inflammatory responses to the model pollutant DEP and that this seems to be related to their reduced capacity to make a cytoprotective Phase II enzymes response. Our studies illuminate why there may be increased susceptibility of certain vulnerable individuals and populations (such as children) to oxidant pollutants and suggest that increasing the body's Phase II responses either by therapeutic or dietary means may counteract this effect. Moreover, the discovery that *in vitro* GST expression is associated with *in vivo* inflammatory responses provides the potential to develop a diagnostic test for susceptibility to oxidant pollutants.

RP.2.4 PUBLICATIONS RESULTING FROM THE CHILDREN'S CENTER (2004–2009)

Publications resulting from collaborations between projects are starred with an asterisk.

- Berhane, K., W. Gauderman, D. Stram and D. Thomas (2004). Statistical issues in studies of the long-term effects of air pollution: The Southern California Children's Health Study. *Stat Sci* 19(3): 414-449.
- Conti, D. V. and W. J. Gauderman (2004). SNPs, haplotypes, and model selection in a candidate gene region: the SIMPLe analysis for multilocus data. *Genet Epidemiol* 27(4): 429-41.
- *Gauderman, W. J., E. Avol, F. Gilliland, H. Vora, D. Thomas, K. Berhane, R. McConnell, N. Kuenzli, F. Lurmann, E. Rappaport, H. Margolis, D. Bates and J. Peters (2004). The effect of air pollution on lung development from 10 to 18 years of age. *N Engl J Med* 351(11): 1057-67.
- *Gilliland, F. D., T. Islam, K. Berhane, R. McConnell and J. Peters (2004). Smoking and asthma incidence in school-aged children: Results from the Children's Health Study. *Am J Respir Crit Care Med* 169(7): A709.
- *Gilliland, F. D., Y.-F. Li, A. Saxon and D. Diaz-Sanchez (2004). Glutathione-S-Transferase M1 and P1 genotypes protect against xenobiotic enhancement of allergic responses. *Lancet* 363: 119-25.
- *Hricko, A. M. (2004). Problems along the new "silk road". *Environ Health Perspect* 112(15): A879 PMID: 1247631.
- Jerrett, M., A. Arain, P. Kanaroglou, B. Beckerman, D. Potoglou, T. Sahsuvaroglu, J. Morrison and C. Giovis (2004). A review and evaluation of intra-urban air pollution exposure models. *J Expo Anal Environ Epidemiol* 15: 185-204.
- Li, Y.-F., Y.-H. Tsao, M. R. Wenten, L. Dubeau and F. D. Gilliland (2004). Intercellular adhesion molecule 1 haplotypes and childhood asthma risk. *Am J Respir Crit Care Med* 169(7): A583.
- *Linn, W. S., M. Avila and H. Gong (2004). Exhaled nitric oxide: sources of error in offline measurement. *Arch Env Health* 59(7): 1-7.
- *Millstein, J., F. Gilliland, K. Berhane, W. J. Gauderman, R. McConnell, E. Avol, E. B. Rappaport and J. M. Peters (2004). Effects of ambient air pollutants on asthma medication use and wheezing among fourth-grade school children from 12 Southern California communities enrolled in The Children's Health Study. *Arch Environ Health* 59(10): 505-14.
- Mueller-Anneling, L., E. Avol, J. M. Peters and P. S. Thorne (2004). Ambient endotoxin concentrations in PM10 from Southern California. *Environ Health Perspect* 112(5): 583-8.
- Salam, M. T., F. Gilliland, Y.-F. Li and B. Langholz (2004). Early life environmental risk factors for asthma: Findings from the Children's Health Study. *Environ Health Perspect* 112(6).
- Thomas, D. C., R. Xie and M. Gebregziabher (2004). Two-stage sampling designs for gene association studies. *Genet Epidemiol* 27: 401-414.
- Conti, D. V., D. O. Stram, J. Molitor, P. Marjoram and D. C. Thomas (2005). Spatial modeling of multilocus data. *Handbook of Statistics in Clinical Oncology* (2nd Ed). In, J. Crowley (Ed), Cancer Research and Biostatistics. 27: 491-507.
- *Diaz-Sanchez, D. and M. Riedl (2005). Diesel effects on human health: a question of stress? *Am J Physiol Lung Cell Mol Physiol* 289(5): L722-3.
- *Gauderman, W. J., E. Avol, F. Lurmann, N. Kuenzli, F. Gilliland, J. Peters and R. McConnell (2005). Childhood asthma and exposure to traffic and nitrogen dioxide. *Epidemiology* 16(6): 737-743.

- Gauderman, W. J. and D. V. Conti (2005). Commentary: Models for longitudinal family data. *Int J Epidemiol* 34(5): 1077-9.
- *Gilliland, F., E. Avol, P. Kinney, M. Jerrett, T. Dvorchak, F. Lurmann, T. Buckley, P. Breysse, G. Keeler, T. de Villiers and R. McConnell (2005). Air pollution exposure assessment for epidemiologic studies of pregnant women and children: lessons learned from the Centers for Children's Environmental Health and Disease Prevention Research. *Environ Health Perspect* 113(10): 1447-54.
- Jerrett, M., A. Arain, P. Kanaroglou, B. Beckerman, D. Potoglou, T. Sahsuvaroglu, J. Morrison and C. Giovis (2005). A review and evaluation of intraurban air pollution exposure models. *J Expo Anal Environ Epidemiol* 15(2): 185-204.
- Kanaroglou, P., M. Jerrett, J. Morrison, B. Beckerman, A. Arain, N. Gilbert and J. Brook (2005). Establishing an air pollution monitoring network for intra-urban population exposure assessment: a location-allocation approach. *Atmos Environ* 39: 2399-2409.
- Li, Y. F., P. C. Lin, L. Dubeau and F. D. Gilliland (2005). The association of clara cell secretory protein gene with childhood asthma and wheezing. *Proc Am Thoracic Soc 2*(Abstract Issue): A619.
- *Li, Y. F., Y. H. Tsao, W. J. Gauderman, D. V. Conti, E. Avol, L. Dubeau and F. D. Gilliland (2005). Intercellular adhesion molecule-1 and childhood asthma. *Hum Genet* 117(5): 476-84.
- Li, Y.-F., B. Langholz, M. T. Salam and F. D. Gilliland (2005). Maternal and grandmaternal smoking patterns are associated with early childhood asthma. *Chest* 127: 1232-1241.
- *Linn, W. S., M. Avila, K. J. Hudson, H. Gong and F. D. Gilliland (2005). Validation of off-site, off-line measurement of exhaled nitric oxide (eNO). *Proc Am Thoracic Soc 2*(Abstract Issue): A309.
- McConnell, R., J. Milam, M. Jerrett, L. Yao and J. Richardson (2005). Parental stress and incident wheeze in a cohort of children from southern California. *Proc Am Thoracic Soc 2*(Abstract Issue): A605.
- McConnell, R., L. Yao, F. Lurmann, N. Kunzli and M. Jerrett (2005). Association of childhood asthma with residence near a major road. *Proc Am Thoracic Soc 2*(Abstract Issue): A522.
- Millstein, J., K. Siegmund, D. V. Conti and W. J. Gauderman (2005). Identifying susceptibility genes by using joint tests of association and linkage and accounting for epistasis. *BMC Genetics* 6(Suppl 1): S147.
- Millstein, J., K. D. Siegmund, D. V. Conti and W. J. Gauderman (2005). Testing association and linkage using affected-sib-parent study designs. *Genet Epidemiol* 29(3): 225-233.
- *Riedl, M. and D. Diaz-Sanchez (2005). Biology of diesel exhaust effects on respiratory function. *J Allergy Clin Immunol* 115(2): 221-8; quiz 229.
- Riedl, M. A., E. M. Landaw, A. Saxon and D. Diaz-Sanchez (2005). Initial high-dose nasal allergen exposure prevents allergic sensitization to a neoantigen. *J Immunol* 174(11): 7440-5.
- Rondeau, V., K. Berhane and D. Thomas (2005). A three-level model for binary time-series data: the effects of air pollution on school absences in the southern California Children's Health Study. *Stat Med* 24(7):1103-15.
- Salam, M. T., B. Langholz, Y.-F. Li and F. D. Gilliland (2005). Maternal fish consumption during pregnancy and risk of childhood asthma. *Journal of Asthma* 42(6): 513-518.
- *Salam, M. T., J. Millstein, Y. F. Li, F. W. Lurmann, H. G. Margolis and F. D. Gilliland (2005). Birth outcomes and prenatal exposure to ozone, carbon monoxide, and particulate matter: results from the Children's Health Study. *Environ Health Perspect* 113(11): 1638-44.
- *Saxon, A. and D. Diaz-Sanchez (2005). Air pollution and allergy: you are what you breathe. *Nat Immunol* 6(3): 223-6.
- Stram, D. O. (2005). Designs for studies of personal exposure to air pollution and the impact of measurement error. *J Toxicol Environ Health A* 68(13-14): 1181-7.
- Stram, D. O. (2005). Software for tag single nucleotide polymorphism selection. *Hum Genomics* 2(2): 144-51.
- Thomas, D. (2005). Why do estimates of the acute and chronic effects of air pollution on mortality differ? *J Toxicol Environ Health A* 68(13-14): 1167-74.
- Thomas, D. C. (2005). The need for a systematic approach to complex pathways in molecular epidemiology. *Cancer Epidemiol Biomarkers Prev* 14(3): 557-9.
- Thomas, D. C., R. W. Haile and D. Duggan (2005). Recent developments in genomewide association scans: a workshop summary and review. *Am J Hum Genet* 77(3): 337-45.
- *Thorne, P. S., N. Metwali, E. Avol and R. S. McConnell (2005). Surface sampling for endotoxin assessment using electrostatic wiping cloths. *Ann Occup Hyg* 49(5): 401-6.
- *Wenten, M., K. Berhane, E. B. Rappaport, E. Avol, W. W. Tsai, W. J. Gauderman, R. McConnell, L. Dubeau and F. D. Gilliland (2005). TNF-308 modifies the effect of second-hand smoke on respiratory illness-related school absences. *Am J Respir Crit Care Med* 172: 1563-1568.

- Wu, J., T. Funk, F. Lurmann and A. Winer (2005). Improving spatial accuracy of roadway networks and geocoded addresses. *Transactions in GIS* 9(4): 585-601.
- *Wu, J., F. Lurmann, A. Winer, R. Lu, R. Turco and T. H. Funk (2005). Development of an individual exposure model for application to the Southern California Children's Health Study. *Atmos Environ* 39(2): 259-273.
- *Diaz-Sanchez, D., R. Rumold and H. Gong, Jr. (2006). Challenge with environmental tobacco smoke exacerbates allergic airway disease in human beings. *J Allergy Clin Immunol* 118(2): 441-6.
- Gauderman, W. J. (2006). Air pollution and children: An unhealthy mix. *N Engl J Med* 355: 78-79.
- *Gilliland, F. D., T. Islam, K. Berhane, W. J. Gauderman, R. McConnell, E. Avol and J. M. Peters (2006). Regular smoking and asthma incidence in adolescents. *Am J Respir Crit Care Med* 174(10): 1094-100.
- *Gilliland, F. D., Y.F. Li, H. Gong, and D. Diaz-Sanchez (2006). Glutathione S-Transferases M1 and P1 prevent aggravation of allergic responses by secondhand smoke. *Am J Respir Crit Care Med* 174(12):1335-41.
- *Gilliland, F. D., W. S. Linn, E. B. Rappaport, E. L. Avol and R. S. McConnell (2006). Exhaled nitric oxide in a population-based sample of school-aged children. *Proc Am Thoracic Soc* 3: A46.
- *Hricko, A. M. (2006). Ships, trucks, and trains: effects of goods movement on environmental health. *Environ Health Perspect* 114(4): A204-5 PMID: 1440794.
- *Islam, T., J. W. Gauderman, K. Berhane, R. McConnell, E. Avol, J. Peters and F. D. Gilliland (2006). Low lung function predicts new onset asthma among children. *Proc Am Thoracic Soc* 3: A153.
- *Jerrett, M., K. Shankardass, J. Gauderman, K. Berhane, N. Kunzli, F. Lurmann, E. Avol, J. Peters, F. Gilliland, J. Molitor, J. Molitor and R. McConnell (2006). Air pollution and asthma incidence in children. *Proc Am Thoracic Soc* 3: A23.
- *Li, Y.-F., W. J. Gauderman, E. Avol, L. Dubeau and F. D. Gilliland (2006). Associations of tumor necrosis factor G-308A with childhood asthma and wheezing. *Am J Respir Crit Care Med* 173(9): 970-6.
- *Linn, W. S., K. R. Anderson, M. Avila, H. Gong, E. B. Rappaport, E. L. Avol and F. D. Gilliland (2006). A survey of exhaled nitric oxide (eNO) in southern California schoolchildren: methodology and data quality. *Proc Am Thoracic Soc* 3: A45.
- *McConnell, R., K. Berhane, J. Molitor, F. Gilliland, N. Künzli, P. Thorne, D. Thomas, J. Gauderman, E. Avol, F. Lurmann, E. Rappaport, M. Jerrett and J. M. Peters (2006). Dog ownership enhances symptomatic responses to air pollution in children with asthma. *Environ Health Perspect* 114(12): 1910-1915 PMID: 1764158.
- *McConnell, R., K. Berhane, L. Yao, M. Jerrett, F. Lurmann, F. Gilliland, N. Kunzli, J. Gauderman, E. Avol, D. Thomas and J. Peters (2006). Traffic, susceptibility, and childhood asthma. *Environ Health Perspect* 114(5): 766-72.
- *McConnell, R., K. Berhane, L. Yao, F. W. Lurmann, E. Avol and J. M. Peters (2006). Predicting residential ozone deficits from nearby traffic. *Sci Total Environ* 363(1-3): 166-74.
- *Millstein, J., D. V. Conti, F. D. Gilliland and W. J. Gauderman (2006). A testing framework for identifying susceptibility genes in the presence of epistasis. *Am J Hum Genet* 78(1): 15-27.
- *Molitor, J., N.-T. Molitor, M. Jerrett, R. McConnell, W. J. Gauderman, K. Berhane and D. Thomas (2006). Bayesian modeling of air pollution health effects with missing exposure data. *Am J Epidemiol* 164(1): 69-76.
- Ross, Z., P. B. English, R. Scalf, R. Gunier, S. Smorodinsky, S. Wall and M. Jerrett (2006). Nitrogen dioxide prediction in Southern California using land use regression modeling: potential for environmental health analyses. *J Expo Sci Environ Epidemiol* 16(2): 106-14.
- *Salam, M. T., P. C. Lin, W. J. Gauderman and F. Gilliland (2006). Genetic polymorphisms in microsomal epoxide hydrolase and childhood asthma. *Proc Am Thoracic Soc* 3: A349.
- *Salam, M. T., H. G. Margolis, R. McConnell, J. A. McGregor, E. L. Avol and F. D. Gilliland (2006). Mode of delivery is associated with asthma and allergy occurrences in children. *Ann Epidemiol* 16(5): 341-6.
- Shankardass, K., R. S. McConnell, J. Milam, Z. Tatalovich, J. P. Wilson and M. Jerrett (2006). The association between community socioeconomic factors and prevalent asthma in southern California school children. *Proc Am Thoracic Soc* 3: A153.
- Wan, J. and D. Diaz-Sanchez (2006). Phase II enzymes induction blocks the enhanced IgE production in B cells by diesel exhaust particles. *J Immunol* 177(5): 3477-83.
- *de Villiers, T., F. Gilliland, Y. Li and D. Diaz-Sanchez (2007). Secondhand smoke, inflammatory responses and Glutathione S-Transferase M1: Results of a randomized challenge experiment. *Epidemiology* 18(5): S39.
- Gauderman, W. J., C. Murcray, F. Gilliland and D. V. Conti (2007). Testing association between disease and multiple SNPs in a candidate gene. *Genet Epidemiol* 31(5): 383-95.

- *Gauderman, W. J., H. Vora, R. McConnell, K. Berhane, F. Gilliland, D. Thomas, F. Lurmann, E. Avol, N. Kunzli, M. Jerrett and J. Peters (2007). Effect of exposure to traffic on lung development from 10 to 18 years of age: a cohort study. *Lancet* 369(9561): 571-7.
- *Islam, T., W. J. Gauderman, K. Berhane, R. McConnell, E. Avol, J. M. Peters and F. D. Gilliland (2007). Relationship between air pollution, lung function and asthma in adolescents. *Thorax* 62(11): 957-63.
- *Islam, T., W. J. Gauderman, R. McConnell and F. D. Gilliland (2007). Heme oxygenase, ozone and incident asthma. *Proc Am Thoracic Soc* 175: A833.
- Kraft, P., Y. C. Yen, D. O. Stram, J. Morrison and W. J. Gauderman (2007). Exploiting gene-environment interaction to detect genetic associations. *Hum Hered* 63(2): 111-9.
- *McConnell, R. (2007). Childhood incident asthma and traffic-related air pollution in a longitudinal cohort study. *Epidemiology* 18(5): S187.
- *McConnell, R., T. Islam, K. Berhane, F. Lurmann, L. Yao, M. Jerrett, J. Gauderman, F. Gilliland, E. Avol, D. Thomas and J. Peters (2007). Childhood incident asthma and traffic-related pollution in a longitudinal cohort study. *Proc Am Thoracic Soc* 175: A304.
- *Molitor, J., M. Jerrett, C. C. Chang, N. T. Molitor, J. Gauderman, K. Berhane, R. McConnell, F. Lurmann, J. Wu, A. Winer and D. Thomas (2007). Assessing uncertainty in spatial exposure models for air pollution health effects assessment. *Environ Health Perspect* 115(8): 1147-53.
- Ritz, S. A., J. Wan and D. Diaz-Sanchez (2007). Sulforaphane-stimulated phase II enzyme induction inhibits cytokine production by airway epithelial cells stimulated with diesel extract. *Am J Physiol Lung Cell Mol Physiol* 292(1): L33-9.
- *Salam, M. T., W. J. Gauderman, R. McConnell, P. C. Lin, M. Wenten and F. D. Gilliland (2007). Transforming growth factor-b1, traffic and childhood asthma. *Proc Am Thoracic Soc* 175: A259.
- Salam, M. T. and F. D. Gilliland (2007). Dietary fat and antioxidant vitamins and respiratory symptoms in children. *Proc Am Thoracic Soc* 175: A272.
- *Salam, M. T., P. C. Lin, E. L. Avol, W. J. Gauderman and F. D. Gilliland (2007). Microsomal epoxide hydrolase, glutathione S-transferase P1, traffic and childhood asthma. *Thorax* 62(12): 1050-7.
- *Salam, M. T., W. J. Gauderman, R. McConnell, P. C. Lin and F. D. Gilliland (2007). Transforming growth factor- 1 C-509T polymorphism, oxidant stress, and early-onset childhood asthma. *Am J Respir Crit Care Med* 176(12): 1192-9.
- *Shankardass, K., R. S. McConnell, J. Milam, K. Berhane, Z. Tatalovich, J. P. Wilson and M. Jerrett (2007). The association between contextual socioeconomic factors and prevalent asthma in a cohort of Southern California school children. *Soc Sci Med* 65(8): 1792-806.
- Thomas, D. C., J. S. Witte and S. Greenland (2007). Dissecting effects of complex mixtures: who's afraid of informative priors? *Epidemiology* 18(2): 186-90.
- Wan, J. and D. Diaz-Sanchez (2007). Antioxidant enzyme induction: a new protective approach against the adverse effects of diesel exhaust particles. *Inhal Toxicol* 19 Suppl 1: 177-82.
- *Avol, E. (2008). The environmental health perspective: a closer look at ports. *Epidemiology* 19(6): S34-S35.
- *Berg, J., R. McConnell, J. Milam, J. Galvan, J. Kotlerman, P. Thorne, C. Jones, R. Ferdman, P. Eggleston, C. Rand, M. A. Lewis, J. Peters and J. Richardson (2008). Rodent allergen in Los Angeles inner city homes of children with asthma. *J Urban Health* 85(1): 52-61 PMID: 2430140.
- *Berhane, K. (2008). Early diagnosis of asthma modifies the effect of air pollution on lung development in children. *Epidemiology* 19(6): S58.
- *Cozen, W., E. Avol, D. Diaz-Sanchez, R. McConnell, W. J. Gauderman, M. G. Cockburn, J. Zadnick, M. Jyrala and T. M. Mack (2008). Use of an electrostatic dust cloth for self-administered home allergen collection. *Twin Res Hum Genet* 11(2): 150-5 PMID: 18361715.
- *Gauderman, J. (2008). A genome wide search for genetic variants that modify air pollution effects on respiratory health. *Epidemiology* 19(6): S58.
- *Gilliland, F. (2008). The role of genetic susceptibility to air pollutants in the etiology of asthma. *Epidemiology* 19(6): S58.
- *Gilliland, F. (2008). Genetic and environmental determinants of global and CpG methylation in children's respiratory system. *Epidemiology* 19(6): S33.
- *Hricko, A. (2008). Global trade comes home: the impacts of international trade and good movement. *Epidemiology* 19(6): S34.
- *Hricko, A. (2008). Global trade comes home: community impacts of goods movement. *Environ Health Perspect* 116(2): A78-81 PMID: 2235209.

- *Hricko, A. and E. Green (2008). Community environmental health assessment: researchers addressing "real world" problems. *Epidemiology* 19(6): S61-S62.
- *Islam, T., R. McConnell, W. J. Gauderman, E. Avol, J. M. Peters and F. D. Gilliland (2008). Ozone, oxidant defense genes, and risk of asthma during adolescence. *Am J Respir Crit Care Med* 177(4): 388-95 PMID: 18048809.
- *Jerrett, M., K. Shankardass, K. Berhane, W. J. Gauderman, N. Kunzli, E. Avol, F. Gilliland, F. Lurmann, J. N. Molitor, J. T. Molitor, D. C. Thomas, J. Peters and R. McConnell (2008). Traffic-related air pollution and asthma onset in children: a prospective cohort study with individual exposure measurement. *Environ Health Perspect* 116(10): 1433-8 PMID: 2569108.
- Kozawa, K., S. Fruin and A. Winer (2008). Using a mobile monitoring platform to characterize pollution concentrations on and near heavily-traveled roadways in communities adjacent to the ports of Los Angeles and Long Beach. *Epidemiology* 19(6): S224-S225.
- *Kunzli, N., L. Perez, F. Lurmann, A. Hricko, B. Penfold and R. McConnell (2008). An attributable risk model for exposures assumed to cause both chronic disease and its exacerbations. *Epidemiology* 19(2): 179-85.
- Lee, Y. L., F. D. Gilliland, J. Y. Wang, Y. C. Lee and Y. L. Guo (2008). Associations of FcepsilonRIbeta E237G polymorphism with wheezing in Taiwanese schoolchildren. *Clin Exp Allergy* 38(3): 413-20 PMID: 18269668.
- *Li, Y., Shah H, D. Diaz-Sanchez and F. Gilliland (2008). GSTM1 genotypes modify the DEP enhancement of nasal responses after cat-allergen challenge. *Epidemiology* 19(6): S124.
- *Li, Y. F., W. J. Gauderman, D. V. Conti, P. C. Lin, E. Avol and F. D. Gilliland (2008). Glutathione s-transferase p1, maternal smoking, and asthma in children: a haplotype-based analysis. *Environ Health Perspect* 116(3): 409-15 PMID: 18335111.
- *Linn, W., E. Rappaport, T. de Villiers, E. Avol, K. Berhane and F. Gilliland (2008). Relationships of ambient nitric oxide and exhaled nitric oxide (by either real-time on-line or delayed off-line measurement) in an epidemiologic survey of schoolchildren. *Epidemiology* 19(6): S137-S138.
- *Linn, W. S., K. T. Berhane, E. B. Rappaport, T. M. Bastain, E. L. Avol and F. D. Gilliland (2008). Relationships of online exhaled, offline exhaled, and ambient nitric oxide in an epidemiologic survey of schoolchildren. *J Expo Sci Environ Epidemiol*. 22 October 2008; doi:10.1038/jes.2008.64.
- *Liu, F., R. McConnell and J. Wu (2008). Association of childhood asthma with school commuting time. *Epidemiology* 19(6): S339-S340.
- *McConnell, R. (2008). Asthma and traffic-related exposure at schools, homes and on roadways. *Epidemiology* 19(6): S58.
- *Milam, J., R. McConnell, L. Yao, K. Berhane, M. Jerrett and J. Richardson (2008). Parental stress and childhood wheeze in a prospective cohort study. *J Asthma* 45(4): 319-23.
- *Perez, L., N. Kuenzli, A. Hricko, E. Avol, F. Lurmann and R. McConnell (2008). Air pollution and the related asthma-burden for children in two impacted communities of southern California. *Epidemiology* 19(6): S130.
- *Peters, J. (2008). Synthesis of scientific results and policy implications from the Children's Health Study. *Epidemiology* 19(6): S58.
- Salam, M. T., T. Islam and F. D. Gilliland (2008). Recent evidence for adverse effects of residential proximity to traffic sources on asthma. *Curr Opin Pulm Med* 14: 3-8 PMID: 18043269.
- *Sawant, A. A., D. R. Cocker, 3rd, J. W. Miller, T. Taliaferro, D. Diaz-Sanchez, W. S. Linn, K. W. Clark and H. Gong, Jr. (2008). Generation and characterization of diesel exhaust in a facility for controlled human exposures. *J Air Waste Manag Assoc* 58(6): 829-37.
- *Shankardass, K., M. Jerrett, J. Milam, J. Richardson, K. Berhane and R. McConnell (2008). Effects of the social environment on incident asthma in childhood: associations with school Title I funding and crime in the community. *Epidemiology* 19(6): S239.
- *Wang, C., M. T. Salam, T. Islam, M. Wenten, W. J. Gauderman and F. D. Gilliland (2008). Effects of in utero and childhood tobacco smoke exposure and beta2-adrenergic receptor genotype on childhood asthma and wheezing. *Pediatrics* 122(1): e107-14 PMID: 18558635.
- *Zhang, Y., R. McConnell and K. Berhane (2008). Ethnic difference in the effect of asthma on pulmonary function in children. *Epidemiology* 19(6): S196-S197.
- *Bastain, T. M., E. Rappaport, T. Islam, M. T. Salam, K. Berhane, W. Linn and F. Gilliland (2009). The effect of exhaled nitric oxide on new onset asthma in the southern California Children's Health Study. *Am J Respir Crit Care Med* 179: A4764.

- *Berhane, K., Y. Zhang, W. Linn, E. Rappaport, T. M. Bastain, F. Lurmann and F. Gilliland (2009). The effect of ambient air pollution on exhaled nitric oxide (FeNO) in southern California schoolchildren. *Am J Respir Crit Care Med* 179: A4732.
- *Breton, C., H. Vora, M. Salam, W. Gauderman and F. Gilliland (2009). Genes in glutathione regulation alter the effect of air pollution on lung function growth. *Am J Respir Crit Care Med* 179: A1661.
- *Breton, C. V., H. Vora, M. T. Salam, T. Islam, M. Wenten, W. J. Gauderman, D. Van Den Berg, K. Berhane, J. M. Peters and F. D. Gilliland (2009). Variation in the GST mu locus and tobacco smoke exposure as determinants of childhood lung function. *Am J Respir Crit Care Med* 179(7): 601-7.
- *Gilliland, F. D. (2009). Outdoor air pollution, genetic susceptibility, and asthma management: opportunities for intervention to reduce the burden of asthma. *Pediatrics* 123(3): S168-S173.
- Islam, T., T. M. Bastain, M. T. Salam and F. Gilliland (2009). Genetic variants of cytokine as determinants of exhaled nitric oxide. *Am J Respir Crit Care Med* 179: A3656.
- *Islam, T., M. T. Salam and F. Gilliland (2009). Variants of cytokine genes in new-onset asthma among adolescents. *Am J Respir Crit Care Med* 179: A1672.
- *Islam, T., K. Berhane, R. McConnell, W. J. Gauderman, E. Avol, J. M. Peters and F. D. Gilliland (2009). Glutathione-S-transferase (GST) P1, GSTM1, exercise, ozone and asthma incidence in school children. *Thorax* 64(3): 197-202.
- *Linn, W., E. Rappaport, K. Berhane, T. M. Bastain, S. Howland, N. Realiza, R. Diaz, L. Valencia and F. Gilliland (2009). Extended exhaled nitric oxide testing in children at schools. *Am J Respir Crit Care Med* 179: A4765.
- *McConnell, R., F. Liu, J. Gauderman and K. Berhane (2009). Association of traffic-related pollutants with alternative asthma phenotypes in the Children's Health Study. *Am J Respir Crit Care Med* 179: A4738.
- Murcray, C. E., J. P. Lewinger and W. J. Gauderman (2009). Gene-environment interaction in genome-wide association studies. *Am J Epidemiol* 169(2): 219-26.
- Riedl, M. A., A. Saxon and D. Diaz-Sanchez (2009). Oral sulforaphane increases Phase II antioxidant enzymes in the human upper airway. *Clin Immunol* 130(3): 244-51.
- *Salam, M. T., T. M. Bastain, T. Islam, M. Wenten and F. Gilliland (2009). Variants in genes in 5-lipoxygenase pathway and exhaled nitric oxide concentrations in children. *Am J Respir Crit Care Med* 179: A1671.
- *Salam, M. T., T. Islam, C. Breton and F. Gilliland (2009). Variants in glutathione s-transferase genes and childhood asthma. *Am J Respir Crit Care Med* 179: A3654.
- *Salam, M. T., T. Islam, W. J. Gauderman and F. D. Gilliland (2009). Roles of arginase variants, atopy, and ozone in childhood asthma. *J Allergy Clin Immunol* 123(3): 596-602, 602 e1-8.
- *Wenten, M., M. Salam, T. Islam, T. Bastain and F. Gilliland (2009). The association of common NOS2A haplotypes on fractional exhaled NO is modified by exposure to second-hand smoke. *Am J Respir Crit Care Med* 179: A1668.
- *Linn, W., E. Rappaport, K. Berhane, T. Bastain, E. Avol and F. Gilliland (In Press). Exhaled Nitric Oxide in a Population-Based Study of Southern California Schoolchildren. *Respiratory Research*.
- *Perez, L., N. Kuenzli, E. Avol, A. Hricko, F. Lurmann, E. Nicholas, F. Gilliland, J. Peters and R. McConnell (In Press). Global goods movement and the local burden of childhood asthma in southern California. *AJPH*.
- *McConnell, R. S., T. Islam, K. Shankardass, M. Jerrett, F. Lurmann, F. D. Gilliland, W. J. Gauderman, E. Avol, N. Kuenzli, L. Yao, J. Peters and K. Berhane (In Review). Childhood incident asthma and traffic-related air pollution in a longitudinal cohort study.
- *Salam, M. T., M. Wenten, P. C. Lin, W. Linn, T. Islam, E. Rappaport, T. M. Bastain, K. Berhane and F. Gilliland (In Review). Nitric oxide synthase and arginase variants, secondhand smoke and exhaled nitric oxide in children.

RP.2.5 SUMMARY OF THE KEY SCIENTIFIC CONTRIBUTIONS OF THE CENTER

The key scientific findings have been presented in Section RP.2 Program Accomplishments. These findings clearly demonstrate that ambient air pollution at current levels has clinically important adverse acute and chronic health effects on children that are likely to contribute to a lifetime of sub-optimal health. The role of PM in causing acute effects has emerged as an important topic of research and progress has been made in understanding these effects.⁹⁰⁻⁹¹ Mechanisms appear to involve oxidative/nitrosative stress and inflammatory pathways. New reports suggest that the chronic effects of PM may be greater than the acute effects^{8, 92} and that consideration of microenvironments and physical activity patterns might be critical. Although the chronic effects of PM are likely to be important, few studies have approached the difficult task of investigating the chronic effects of long-term exposures across the life course. The extensive CHS resources, the levels of

exposure in southern California, the new analytic methods and the experience and highly collaborative multidisciplinary team contribute to the high probability that the program of research proposed in this application will fill key gaps in the knowledge base to advance our scientific understanding of respiratory disease in general, and the specific contribution of air pollution to the occurrence of respiratory diseases.

Literature Cited

1. McConnell R, Berhane K, Gilliland F, et al. Asthma in exercising children exposed to ozone: A cohort study. *Lancet* 2002;359:386-91.
2. McConnell R, Berhane K, Gilliland F, et al. Prospective Study of Air Pollution and Bronchitic Symptoms in Children with Asthma. *Am J Respir Crit Care Med* 2003;168:790-7.
3. McConnell R, Berhane K, Yao L, et al. Traffic, susceptibility, and childhood asthma. *Environ Health Perspect* 2006;114:766-72.
4. Gauderman WJ. Traffic-related pollution and lung function growth. *Proceedings of the American Thoracic Society* 2006:A78.
5. Gauderman W, Vora H, McConnell R, et al. The effect of residential exposure to traffic on lung development from 10 to 18 years of age. *Lancet* in press.
6. Gauderman WJ, Avol E, Lurmann F, et al. Childhood asthma and exposure to traffic and nitrogen dioxide. *Epidemiology* 2005;16:737-43.
7. Gilliland F. Residential traffic, asthma and school absence. *Proceedings of the American Thoracic Society* 2006:A78.
8. Gauderman WJ, Avol E, Gilliland F, et al. The effect of air pollution on lung development from 10 to 18 years of age. *N Engl J Med* 2004;351:1057-67.
9. Peters JM, Avol E, Gauderman WJ, et al. A study of twelve Southern California communities with differing levels and types of air pollution. II. Effects on pulmonary function. *Am J Respir Crit Care Med* 1999;159:768-75.
10. Peters JM, Avol E, Navidi W, et al. A study of twelve Southern California communities with differing levels and types of air pollution. I. Prevalence of respiratory morbidity. *Am J Respir Crit Care Med* 1999;159:760-7.
11. Gilliland FD, Berhane K, Rappaport EB, et al. The effects of ambient air pollution on school absenteeism due to respiratory illnesses. *Epidemiology* 2001;12:43-54.
12. Gilliland FD, Li Y-F, Tsai W, Dubeau L, Avol E, Peters JM. TNF α -308 genotype and ozone effects on asthma and wheezing: Results from the Children's Health Study (CHS). *Am J Respir Crit Care Med* 2003;167:A580.
13. Jerrett M, Shankardass K, Gauderman W, et al. Air pollution and asthma incidence in children. *American Journal of Respiratory and Critical Care Medicine* 2006;3:A23.
14. McConnell R, Berhane K, Gilliland F, et al. Air Pollution and Bronchitic Symptoms in Southern California Children with Asthma. *Environ Health Perspect* 1999;107:757-60.
15. Molitor J, Molitor N-T, Jerrett M, et al. Bayesian modeling of air pollution health effects with missing exposure data. *Am J Epidemiol* 2006;164:69-76.
16. Gilliland FD, McConnell R, Peters J, Gong H, Jr. A theoretical basis for investigating ambient air pollution and children's respiratory health. *Environ Health Perspect* 1999;107 Suppl 3:403-7.
17. Gilliland FD, Li Y-F, Saxon A, Diaz-Sanchez D. Glutathione-S-Transferase M1 and P1 genotypes protect against xenobiotic enhancement of allergic responses. *Lancet* 2004;363:119-25.
18. Molfino NA, Jeffery PK. Chronic obstructive pulmonary disease: Histopathology, inflammation and potential therapies. *Pulm Pharmacol Ther* 2006.
19. Rahman I. Oxidative stress in pathogenesis of chronic obstructive pulmonary disease: cellular and molecular mechanisms. *Cell Biochem Biophys* 2005;43:167-88.
20. Chaudhuri N, Dower SK, Whyte MK, Sabroe I. Toll-like receptors and chronic lung disease. *Clin Sci (Lond)* 2005;109:125-33.

21. Psarras S, Caramori G, Contoli M, Papadopoulos N, Papi A. Oxidants in asthma and in chronic obstructive pulmonary disease (COPD). *Curr Pharm Des* 2005;11:2053-62.
22. van der Vliet A, Cross CE. Oxidants, nitrosants, and the lung. *American Journal of Medicine* 2000;109:398-421.
23. Wan J, Diaz-Sanchez D. Phase II enzymes induction blocks the enhanced IgE production in B cells by diesel exhaust particles. *J Immunol* 2006;177:3477-83.
24. Ritz SA, Wan J, Diaz-Sanchez D. Sulforaphane-stimulated phase II enzyme induction inhibits cytokine production by airway epithelial cells stimulated with diesel extract. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L33-9.
25. Gauderman WJ, Avol E, Lurmann F, et al. Childhood asthma and exposure to traffic and nitrogen dioxide. *Epidemiology* 2005;16:737-43.
26. Gauderman WJ, Vora H, McConnell R, et al. Effect of exposure to traffic on lung development from 10 to 18 years of age: a cohort study. *Lancet* 2007;369:571-7.
27. McConnell R, Liu F, Gauderman J, Berhane K. Association of traffic-related pollutants with alternative asthma phenotypes in the Children's Health Study. *Am J Respir Crit Care Med* 2009;179:A4738.
28. McConnell R, Islam T, Berhane K, et al. Childhood incident asthma and traffic-related pollution in a longitudinal cohort study. *Proc Am Thoracic Soc* 2007;175:A304.
29. McConnell R. Asthma and traffic-related exposure at schools, homes and on roadways. *Epidemiology* 2008;19:S58.
30. Jerrett M, Shankardass K, Gauderman J, et al. Air pollution and asthma incidence in children. *Proc Am Thoracic Soc* 2006;3:A23.
31. Jerrett M, Shankardass K, Berhane K, et al. Traffic-related air pollution and asthma onset in children: a prospective cohort study with individual exposure measurement. *Environ Health Perspect* 2008;116:1433-8.
32. Liu F, McConnell R, Wu J. Association of Childhood Asthma with School Commuting Time. *Epidemiology* 2008;19:S339-S40.
33. Li Y-F, Langholz B, Salam MT, Gilliland FD. Maternal and grandmaternal smoking patterns are associated with early childhood asthma. *Chest* 2005;127:1232-41.
34. Gilliland FD, Islam T, Berhane K, et al. Regular smoking and asthma incidence in adolescents. *Am J Respir Crit Care Med* 2006;174:1094-100.
35. Salam MT, Islam T, Gauderman WJ, Gilliland FD. Roles of arginase variants, atopy, and ozone in childhood asthma. *J Allergy Clin Immunol* 2009;123:596-602, e1-8.
36. Wenten M, Salam M, Islam T, Bastain T, Gilliland F. The association of common NOS2A haplotypes on fractional exhaled NO is modified by exposure to second-hand smoke. *Am J Respir Crit Care Med* 2009;179:A1668.
37. Salam MT, Wenten M, Lin PC, et al. Nitric oxide synthase and arginase variants, secondhand smoke and exhaled nitric oxide in children. Submitted.
38. Salam MT, Bastain TM, Islam T, Wenten M, Gilliland F. Variants in genes in 5-lipoxygenase pathway and exhaled nitric oxide concentrations in children. *Am J Respir Crit Care Med* 2009;179:A1671.
39. Islam T, Salam MT, Gilliland F. Variants of cytokine genes in new-onset asthma among adolescents. *Am J Respir Crit Care Med* 2009;179:A1672.
40. Islam T, Bastain TM, Salam MT, Gilliland F. Genetic variants of cytokine as determinants of exhaled nitric oxide. *Am J Respir Crit Care Med* 2009;179:A3656.
41. Li Y-F, Gauderman WJ, Avol E, Dubeau L, Gilliland FD. Associations of tumor necrosis factor G-308A with childhood asthma and wheezing. *Am J Respir Crit Care Med* 2006;173:970-6.
42. Wenten M, Berhane K, Rappaport EB, et al. TNF-308 modifies the effect of second-hand smoke on respiratory illness-related school absences. *Am J Respir Crit Care Med* 2005;172:1563-8.
43. Li Y-F, Tsao Y-H, Wenten MR, Dubeau L, Gilliland FD. Intercellular adhesion molecule 1 haplotypes and childhood asthma risk. *Am J Respir Crit Care Med* 2004;169:A583.
44. Li YF, Tsao YH, Gauderman WJ, et al. Intercellular adhesion molecule-1 and childhood asthma. *Hum Genet* 2005;117:476-84.
45. Salam MT, Lin PC, Gauderman WJ, Gilliland F. Genetic polymorphisms in microsomal epoxide hydrolase and childhood asthma. *Proc Am Thoracic Soc* 2006;3:A349.

46. Salam MT, Lin PC, Avol EL, Gauderman WJ, Gilliland FD. Microsomal epoxide hydrolase, glutathione S-transferase P1, traffic and childhood asthma. *Thorax* 2007;62:1050-7.
47. Li YF, Lin PC, Dubeau L, Gilliland FD. The association of clara cell secretory protein gene with childhood asthma and wheezing. *Proceedings of the American Thoracic Society* 2005;2:A619.
48. Breton C, Vora H, Salam M, Gauderman W, Gilliland F. Genes in glutathione regulation alter the effect of air pollution on lung function growth. *Am J Respir Crit Care Med* 2009;179:A1661.
49. Salam MT, Islam T, Breton C, Gilliland F. Variants in glutathione s-transferase genes and childhood asthma. *Am J Respir Crit Care Med* 2009;179:A3654.
50. Islam T, Gauderman JW, Berhane K, et al. Low lung function predicts new onset asthma among children. *Proc Am Thoracic Soc* 2006;3:A153.
51. Islam T, Gauderman WJ, Berhane K, et al. Relationship between air pollution, lung function and asthma in adolescents. *Thorax* 2007;62:957-63.
52. Islam T, Berhane K, McConnell R, et al. Glutathione-S-transferase (GST) P1, GSTM1, exercise, ozone and asthma incidence in school children. *Thorax* 2009;64:197-202.
53. Islam T, Gauderman WJ, McConnell R, Gilliland FD. Heme oxygenase, ozone and incident asthma. *Proc Am Thoracic Soc* 2007;175:A833.
54. Islam T, McConnell R, Gauderman WJ, Avol E, Peters JM, Gilliland FD. Ozone, oxidant defense genes, and risk of asthma during adolescence. *Am J Respir Crit Care Med* 2008;177:388-95.
55. Breton CV, Vora H, Salam MT, et al. Variation in the GST mu locus and tobacco smoke exposure as determinants of childhood lung function. *Am J Respir Crit Care Med* 2009;179:601-7.
56. Li YF, Gauderman WJ, Conti DV, Lin PC, Avol E, Gilliland FD. Glutathione s-transferase p1, maternal smoking, and asthma in children: a haplotype-based analysis. *Environ Health Perspect* 2008;116:409-15.
57. Shankardass K, McConnell RS, Milam J, et al. The association between contextual socioeconomic factors and prevalent asthma in a cohort of Southern California school children. *Soc Sci Med* 2007;65:1792-806.
58. Shankardass K, Jerrett M, Milam J, Richardson J, Berhane K, McConnell R. Effects of the social environment on incident asthma in childhood: associations with school Title I funding and crime in the community. *Epidemiology* 2008;19:S239.
59. Milam J, McConnell R, Yao L, Berhane K, Jerrett M, Richardson J. Parental stress and childhood wheeze in a prospective cohort study. *J Asthma* 2008;45:319-23.
60. Shankardass K, McConnell RS, Milam J, Tatalovich Z, Wilson JP, Jerrett M. The association between community socioeconomic factors and prevalent asthma in southern California school children. *Proc Am Thoracic Soc* 2006;3:A153.
61. Gilliland FD, Linn WS, Rappaport EB, Avol EL, McConnell RS. Exhaled nitric oxide in a population-based sample of school-aged children. *Proc Am Thoracic Soc* 2006;3:A46.
62. Linn WS, Anderson KR, Avila M, et al. A survey of exhaled nitric oxide (eNO) in southern California schoolchildren: methodology and data quality. *Proc Am Thoracic Soc* 2006;3:A45.
63. Linn W, Rappaport E, Berhane K, Bastain T, Avol E, Gilliland F. Exhaled Nitric Oxide in a Population-Based Study of Southern California Schoolchildren. *Respiratory Research* In Press.
64. Berhane K, Zhang Y, Linn W, et al. The effect of ambient air pollution on exhaled nitric oxide (FeNO) in southern California schoolchildren. *Am J Respir Crit Care Med* 2009;179:A4732.
65. Bastain TM, Rappaport E, Islam T, et al. The effect of exhaled nitric oxide on new onset asthma in the southern California Children's Health Study. *Am J Respir Crit Care Med* 2009;179:A4764.
66. Riedl MA, Landaw EM, Saxon A, Diaz-Sanchez D. Initial high-dose nasal allergen exposure prevents allergic sensitization to a neoantigen. *J Immunol* 2005;174:7440-5.
67. Diaz-Sanchez D, Rumold R, Gong H, Jr. Challenge with environmental tobacco smoke exacerbates allergic airway disease in human beings. *J Allergy Clin Immunol* 2006;118:441-6.
68. Wan J, Diaz-Sanchez D. Antioxidant enzyme induction: a new protective approach against the adverse effects of diesel exhaust particles. *Inhal Toxicol* 2007;19 Suppl 1:177-82.
69. Riedl MA, Saxon A, Diaz-Sanchez D. Oral sulforaphane increases Phase II antioxidant enzymes in the human upper airway. *Clin Immunol* 2009;130:244-51.

70. Linn WS, Avila M, Gong H. Exhaled nitric oxide: sources of error in offline measurement. *Arch Env Health* 2004;59:1-7.
71. Linn WS, Avila M, Hudson KJ, Gong H, Gilliland FD. Validation of off-site, off-line measurement of exhaled nitric oxide (eNO). *Proc Am Thoracic Soc* 2005;2:A309.
72. Linn WS, Berhane KT, Rappaport EB, Bastain TM, Avol EL, Gilliland FD. Relationships of online exhaled, offline exhaled, and ambient nitric oxide in an epidemiologic survey of schoolchildren. *J Expo Sci Environ Epidemiol* 2008.
73. Linn W, Rappaport E, Berhane K, et al. Extended exhaled nitric oxide testing in children at schools. *Am J Respir Crit Care Med* 2009;179:A4765.
74. Jerrett M, Arain A, Kanaroglou P, et al. A review and evaluation of intraurban air pollution exposure models. *J Expo Anal Environ Epidemiol* 2005;15:185-204.
75. Kanaroglou P, Jerrett M, Morrison J, et al. Establishing an air pollution monitoring network for intra-urban population exposure assessment: a location-allocation approach. *Atmos Environ* 2005;39:2399-409.
76. McConnell R, Berhane K, Yao L, Lurmann FW, Avol E, Peters JM. Predicting residential ozone deficits from nearby traffic. *Sci Total Environ* 2006;363:166-74.
77. Berhane K, Gauderman W, Stram D, Thomas D. Statistical issues in studies of the long term effects of air pollution: The Southern California Children's Health Study. *Statist Sci* 2004;19:414-49.
78. Millstein J, Conti DV, Gilliland FD, Gauderman WJ. A testing framework for identifying susceptibility genes in the presence of epistasis. *Am J Hum Genet* 2006;78:15-27.
79. Conti DV, Gauderman WJ. SNPs, haplotypes, and model selection in a candidate gene region: the SIMPle analysis for multilocus data. *Genet Epidemiol* 2004;27:429-41.
80. Gauderman WJ, Murcray C, Gilliland F, Conti DV. Testing association between disease and multiple SNPs in a candidate gene. *Genet Epidemiol* 2007;31:383-95.
81. Murcray CE, Lewinger JP, Gauderman WJ. Gene-environment interaction in genome-wide association studies. *Am J Epidemiol* 2009;169:219-26.
82. Kunzli N, Perez L, Lurmann F, Hricko A, Penfold B, McConnell R. An attributable risk model for exposures assumed to cause both chronic disease and its exacerbations. *Epidemiology* 2008;19:179-85.
83. Perez L, Kuenzli N, Hricko A, Avol E, Lurmann F, McConnell R. Air pollution and the related asthma-burden for children in two impacted communities of southern California. *Epidemiology* 2008;19:S130.
84. Perez L, Kuenzli N, Avol E, et al. Global goods movement and the local burden of childhood asthma in southern California. *AJPH In Press*.
85. Hricko AM. Problems along the new "silk road". *Environ Health Perspect* 2004;112:A879.
86. Hricko AM. Ships, trucks, and trains: effects of goods movement on environmental health. *Environ Health Perspect* 2006;114:A204-5.
87. Hricko A. Global trade comes home: the impacts of international trade and good movement. *Epidemiology* 2008;19:S34.
88. Hricko A. Global trade comes home: community impacts of goods movement. *Environ Health Perspect* 2008;116:A78-81.
89. Gilliland FD. Outdoor air pollution, genetic susceptibility, and asthma management: opportunities for intervention to reduce the burden of asthma. *Pediatrics* 2009;123:S168-S73.
90. National Research Council. Research Priorities for Airborne Particulate Matter: IV. Continuing Research Progress. Washington, D.C.: National Academy Press; 2004.
91. Pope CA, 3rd, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 2006;56:709-42.
92. Jerrett M, Burnett RT, Ma R, et al. Spatial Analysis of Air Pollution and Mortality in Los Angeles. *Epidemiology* 2005;16:727-36.
93. Islam T, Breton C, Salam MT, et al. Role of inducible nitric oxide synthase in asthma risk and lung function growth during adolescence. *Thorax* 2010;65:139-45.
94. Wenten M, Gauderman WJ, Berhane K, Lin PC, Peters J, Gilliland FD. Functional variants in the catalase and myeloperoxidase genes, ambient air pollution, and respiratory-related school absences: an example of epistasis in gene-environment interactions. *Am J Epidemiol* 2009;170:1494-501.

95. Perez L, Kunzli N, Avol E, et al. Global goods movement and the local burden of childhood asthma in southern California. *Am J Public Health* 2009;99 Suppl 3:S622-8.
96. Jerrett M, McConnell R, Chang CC, et al. Automobile traffic around the home and attained body mass index: a longitudinal cohort study of children aged 10-18 years. *Prev Med* 2010;50 Suppl 1:S50-8.
97. Linn WS, Rappaport EB, Berhane KT, Bastain TM, Salam MT, Gilliland FD. Extended exhaled nitric oxide analysis in field surveys of schoolchildren: a pilot test. *Pediatr Pulmonol* 2009;44:1033-42.
99. Breton CV, Byun HM, Wenten M, Pan F, Yang A, Gilliland FD. Prenatal tobacco smoke exposure affects global and gene-specific DNA methylation. *Am J Respir Crit Care Med* 2009;180:462-7.
101. Seibold MA, Reese TA, Choudhry S, et al. Differential enzymatic activity of common haplotypic versions of the human acidic Mammalian chitinase protein. *J Biol Chem* 2009;284:19650-8.
102. Himes BE, Hunninghake GM, Baurley JW, et al. Genome-wide association analysis identifies PDE4D as an asthma-susceptibility gene. *Am J Hum Genet* 2009;84:581-93.
103. Wenten M, Li YF, Lin PC, et al. In utero smoke exposure, glutathione S-transferase P1 haplotypes, and respiratory illness-related absence among schoolchildren. *Pediatrics* 2009;123:1344-51.
104. Linn WS, Rappaport EB, Berhane KT, Bastain TM, Avol EL, Gilliland FD. Exhaled nitric oxide in a population-based study of southern California schoolchildren. *Respir Res* 2009;10:28.
105. Gilliland FD. Outdoor air pollution, genetic susceptibility, and asthma management: opportunities for intervention to reduce the burden of asthma. *Pediatrics* 2009;123 Suppl 3:S168-73.
106. Mirabelli MC, Kunzli N, Avol E, et al. Respiratory symptoms following wildfire smoke exposure: airway size as a susceptibility factor. *Epidemiology* 2009;20:451-9.
107. Lee YL, McConnell R, Berhane K, Gilliland FD. Ambient ozone modifies the effect of tumor necrosis factor G-308A on bronchitic symptoms among children with asthma. *Allergy* 2009;64:1342-8.
108. Linn WS, Berhane KT, Rappaport EB, Bastain TM, Avol EL, Gilliland FD. Relationships of online exhaled, offline exhaled, and ambient nitric oxide in an epidemiologic survey of schoolchildren. *J Expo Sci Environ Epidemiol* 2009;19:674-81.
109. Breton CV, Salam MT, Vora H, Gauderman WJ, Gilliland FD. Genetic Variation in the Glutathione Synthesis Pathway, Air Pollution, and Children's Lung Function Growth. *Am J Respir Crit Care Med* 2010.
110. McConnell R, Liu F, Wu J, Lurmann F, Peters J, Berhane K. Asthma and school commuting time. *J Occup Environ Med* 2010;52:827-8.
111. Bastain TM, Islam T, Berhane KT, et al. Exhaled nitric oxide, susceptibility and new-onset asthma in the children's health study. *Eur Respir J* 2010.
112. Hunninghake GM, Soto-Quiros ME, Avila L, et al. TSLP polymorphisms are associated with asthma in a sex-specific fashion. *Allergy* 2010.
114. McConnell R, Islam T, Shankardass K, et al. Childhood incident asthma and traffic-related air pollution at home and school. *Environ Health Perspect* 2010;118:1021-6.
115. Kunzli N, Jerrett M, Garcia-Esteban R, et al. Ambient air pollution and the progression of atherosclerosis in adults. *PLoS One* 2010;5:e9096.
116. Stidley CA, Picchi MA, Leng S, et al. Multivitamins, folate, and green vegetables protect against gene promoter methylation in the aerodigestive tract of smokers. *Cancer Res* 2010;70:568-74.